arXiv:1510.02758v1 [math.RA] 9 Oct 2015

COMMENSURABILITY OF AUTOMORPHISM GROUPS

ALEX BARTEL AND HENDRIK W. LENSTRA JR.

Abstract. We develop a theory of commensurability of groups, of rings, and of modules. It allows us, in certain cases, to compare sizes of automorphism groups of modules, even when those are infinite. This work is motivated by the Cohen–Lenstra heuristics on class groups. The number theoretic implications will be addressed in a separate paper.

1. Introduction

Often, when a mathematical object is drawn in some "random" manner, the probability that it is isomorphic to a given object is inversely proportional to the size of the automorphism group of the latter. The Cohen–Lenstra heuristics [3, 4], which make predictions on the distribution of class groups of "random" algebraic number fields, are, in fact, a special case of this rule, provided that one passes to Arakelov class groups. However, Arakelov class groups may have infinitely many automorphisms, so that a difficulty arises in comparing the sizes of their automorphism groups. This difficulty is addressed in the present paper; the number theoretic implications will appear in a later one.

Our main results, formulated as Theorems 1.1 and 1.2 below, express that, for certain pairs of modules L and M over certain types of ring, one can meaningfully define the ratio of the size of the automorphism group $\operatorname{Aut} M$ of M to the size of $\operatorname{Aut} L$, even when their orders $\#\operatorname{Aut} M$ and $\#\operatorname{Aut} L$ are infinite. If $\operatorname{Aut} L$ can be naturally embedded in $\operatorname{Aut} M$ as a subgroup of finite index, then the ratio mentioned may be defined to be that index. Our approach consists of giving a canonical definition of such an index in a more general situation. We shall write $\operatorname{ia}(L, M)$ for the "index of automorphism groups" that we define.

Denote by $\mathbb{Z}$ the ring of integers, by $\mathbb{Q}$ the field of rational numbers, by $\mathbb{Q}_{>0}$ the multiplicative group of positive rational numbers, by $R[G]$ the group ring of a group G over a ring R, and by $(G : H)$ the index of a subgroup H of a group G. By "module" we shall always mean "left module".

Theorem 1.1. *Let G be a finite group, let V be a finitely generated $\mathbb{Q}[G]$-module, and put $\mathcal{S} = \{L : L$ is a finitely generated $\mathbb{Z}[G]$-module with $\mathbb{Q} \otimes_{\mathbb{Z}} L \cong V$ as $\mathbb{Q}[G]$-modules$\}$. Then there exists a unique function $\operatorname{ia}\colon \mathcal{S} \times \mathcal{S} \to \mathbb{Q}_{>0}$ such that*

(a) *if L, L', M, $M' \in \mathcal{S}$ and $L \cong L'$, $M \cong M'$, then $\operatorname{ia}(L, M) = \operatorname{ia}(L', M')$;*
(b) *if L, M, $N \in \mathcal{S}$, then $\operatorname{ia}(L, M) \cdot \operatorname{ia}(M, N) = \operatorname{ia}(L, N)$;*

Date: October 12, 2015.

The first author gratefully acknowledges the financial support of the Royal Commission for the Exhibition of 1851.

(c) *if $M \in \mathcal{S}$, and $L \subset M$ is a submodule of finite index, then with $H = \{\sigma \in \operatorname{Aut} M : \sigma L = L\}$ and $\rho\colon H \to \operatorname{Aut} L$ mapping $\sigma \in H$ to $\sigma|L$, one has*

$$\operatorname{ia}(L, M) = \frac{(\operatorname{Aut} M : H) \cdot \#\ker \rho}{(\operatorname{Aut} L : \rho H)}.$$

To explain part (c), we remark that it is not hard to show that one has $L \in \mathcal{S}$, and that the three cardinal numbers $(\operatorname{Aut} M : H)$, $\#\ker\rho$, $(\operatorname{Aut} L : \rho H)$ are finite (see Section 7). Since these three numbers may be thought of as the ratio of the sizes of $\operatorname{Aut} M$ and H, of H and ρH, and of $\operatorname{Aut} L$ and ρH, respectively, one may think of the expression in (c) as the ratio of the sizes of $\operatorname{Aut} M$ and $\operatorname{Aut} L$. The same argument shows that one has indeed $\operatorname{ia}(L, M) = (\#\operatorname{Aut} M)/\#\operatorname{Aut} L$ if $\operatorname{Aut} M$ and $\operatorname{Aut} L$ are finite.

As an example, let G be the trivial group, and put $n = \dim_{\mathbb{Q}} V$. Then each $L \in \mathcal{S}$ is isomorphic to the direct sum of $\mathbb{Z}^n$ with a finite abelian group L_0, and $\operatorname{Aut} L$ is isomorphic to a semidirect product $\operatorname{Hom}(\mathbb{Z}^n, L_0) \rtimes (\operatorname{Aut} L_0 \times \operatorname{GL}(n, \mathbb{Z}))$, where both $\operatorname{Hom}(\mathbb{Z}^n, L_0)$ and $\operatorname{Aut} L_0$ are finite. Writing $M \in \mathcal{S}$ similarly, and "cancelling" $\operatorname{GL}(n, \mathbb{Z})$, one is led to believe that

$$\operatorname{ia}(L, M) = \frac{\#\operatorname{Hom}(\mathbb{Z}^n, M_0) \cdot \#\operatorname{Aut} M_0}{\#\operatorname{Hom}(\mathbb{Z}^n, L_0) \cdot \#\operatorname{Aut} L_0} = \frac{(\#M_0)^n \cdot \#\operatorname{Aut} M_0}{(\#L_0)^n \cdot \#\operatorname{Aut} L_0}.$$

Making this informal argument rigorous (see Proposition 8.5), one discovers that if a function as in Theorem 1.1 exists, it must be given by the formula just stated. However, that this formula does define a function meeting all conditions, in particular (c), is not obvious. Likewise, for general G the uniqueness statement of Theorem 1.1 is easy by comparison to the existence statement. Our proof of Theorem 1.1 is given in Section 8.

There is little doubt that one can prove Theorem 1.1 using a suitable theory of covolumes of arithmetic groups. Instead, we will give an entirely algebraic proof, obtaining the theorem as a special case of a much more general result, of which the formulation requires some terminological preparation.

Isogenies. A *group isogeny* is a group homomorphism $f\colon H \to G$ with $\#\ker f < \infty$ and $(G : fH) < \infty$, and its *index* $\mathrm{i}(f)$ is defined to be $(G : fH)/\#\ker f$. For a ring R, an *R-module isogeny* is an R-module homomorphism that is an isogeny as a map of additive groups. A *ring isogeny* is a ring homomorphism that is an isogeny as a map of additive groups. The index of an isogeny of one of the latter two types is the index of the induced group isogeny on the additive groups.

Commensurabilities. If X, Y are objects of a category $\mathcal{C}$, then a *correspondence* from X to Y in $\mathcal{C}$ is a triple $c = (W, f, g)$, where W is an object of $\mathcal{C}$ and $f\colon W \to X$ and $g\colon W \to Y$ are morphisms in $\mathcal{C}$; we will often write $c\colon X \rightleftharpoons Y$ to indicate a correspondence. A *group commensurability* is a correspondence $c = (W, f, g)$ in the category of groups for which both f and g are isogenies, and the *index* $\mathrm{i}(c)$ of such an isogeny is defined to be $\mathrm{i}(g)/\mathrm{i}(f)$. For a ring R, one defines R-module commensurabilities and their indices analogously, replacing the category of groups by the category of R-modules. Likewise, one defines ring commensurabilities and their indices.

Endomorphisms and automorphisms. Let R be a ring, and let $c = (N, f, g)\colon L \rightleftharpoons M$ be a correspondence of R-modules. We define the *endomorphism*

ring $\operatorname{End} c$ of c to be the subring $\{(\lambda, \nu, \mu) \in (\operatorname{End} L) \times (\operatorname{End} N) \times (\operatorname{End} M) : \lambda f = f\nu, \mu g = g\nu\}$ of the product ring $(\operatorname{End} L) \times (\operatorname{End} N) \times (\operatorname{End} M)$. There are natural ring homomorphisms $\operatorname{End} c \to \operatorname{End} L$ and $\operatorname{End} c \to \operatorname{End} M$ sending (λ, ν, μ) to λ and μ, respectively; we shall write $\mathrm{e}(c)\colon \operatorname{End} L \rightleftharpoons \operatorname{End} M$ for the ring correspondence consisting of $\operatorname{End} c$ and those two ring homomorphisms. Similarly, writing $E^\times$ for the multiplicative group of invertible elements of a ring E, we define the *automorphism group* $\operatorname{Aut} c$ of c to be the group $(\operatorname{End} c)^\times$, and we write $\mathrm{a}(c)\colon \operatorname{Aut} L \rightleftharpoons \operatorname{Aut} M$ for the group correspondence consisting of $\operatorname{Aut} c$ and the natural maps $\operatorname{Aut} c \to \operatorname{Aut} L$, $\operatorname{Aut} c \to \operatorname{Aut} M$.

A *domain* is a commutative ring without zero divisors and with $1 \neq 0$. A ring is *semisimple* if all short exact sequences of modules over the ring split.

We can now formulate the general result that we announced.

Theorem 1.2. *Let Z be an infinite domain such that for all non-zero $m \in Z$ the ring Z/mZ is finite, let Q be the field of fractions of Z, let A be a semisimple Q-algebra of finite vector space dimension over Q, let $R \subset A$ be a sub-Z-algebra with $Q \cdot R = A$, and let L, M be finitely generated R-modules. Then:*

(a) *there is an R-module commensurability $L \rightleftharpoons M$ if and only if the A-modules $Q \otimes_Z L$ and $Q \otimes_Z M$ are isomorphic;*
(b) *if $c\colon L \rightleftharpoons M$ is an R-module commensurability, then $\mathrm{e}(c)\colon \operatorname{End} L \rightleftharpoons \operatorname{End} M$ is a ring commensurability, and $\mathrm{a}(c)\colon \operatorname{Aut} L \rightleftharpoons \operatorname{Aut} M$ is a group commensurability;*
(c) *if c, $c'\colon L \rightleftharpoons M$ are R-module commensurabilities, then one has*

$$\mathrm{i}(\mathrm{e}(c)) = \mathrm{i}(\mathrm{e}(c')), \qquad \mathrm{i}(\mathrm{a}(c)) = \mathrm{i}(\mathrm{a}(c')).$$

Part (c) is the essential statement of Theorem 1.2. It shows that one can define $\mathrm{ia}(L, M) = \mathrm{i}(\mathrm{a}(c))$, independently of c, if one has $Q \otimes_Z L \cong_A Q \otimes_Z M$ and $c\colon L \rightleftharpoons M$ is an R-module commensurability. One deduces the existence part of Theorem 1.1 from Theorem 1.2 by putting $Z = \mathbb{Z}$, $Q = \mathbb{Q}$, $A = \mathbb{Q}[G]$, and $R = \mathbb{Z}[G]$.

We prove Theorem 1.2(a) and (b) as Theorems 6.3 and 7.2, respectively, but with the semisimplicity assumption on A replaced by the condition that R be left-noetherian; the following result shows that this is a weaker condition.

Theorem 1.3. *Let R be a ring as in the statement of Theorem 1.2. Then R is left-noetherian and right-noetherian.*

For a proof, see Theorem 4.2. The point of Theorem 1.3 is that R is not required to be finitely generated as a Z-module. As an aside, we will characterise, in Theorem 4.5, the rings Z satisfying the hypotheses of Theorem 1.2.

The full statement of Theorem 1.2 will be proven in Section 8. There we also give an example (Example 8.2) showing that Theorem 1.2(c) may fail if R is left-noetherian, but A is not semisimple.

Isogenies, commensurabilities, and their indices have many formal properties, and it is to these that Section 2 is devoted. Among other things, we shall, under certain conditions, define the *composition* $d \circ c$ of two correspondences d and c. Composition plays an important role in the proof of

Theorem 1.2. In fact, the index of a commensurability is multiplicative in compositions of commensurabilities. More importantly, and less formally, if in the situation of Theorem 1.2 one has commensurabilities $c\colon L \rightleftharpoons M$ and $d\colon M \rightleftharpoons N$ of finitely generated R-modules, then one has

$$\mathrm{i}(\mathrm{e}(d \circ c)) = \mathrm{i}(\mathrm{e}(d))\mathrm{i}(\mathrm{e}(c)), \qquad \mathrm{i}(\mathrm{a}(d \circ c)) = \mathrm{i}(\mathrm{a}(d))\mathrm{i}(\mathrm{a}(c))$$

(see Theorem 7.3). This property is the first key to the proof of Theorem 1.2(c).

The following elegant result, which will be proven as Theorem 3.8, will be useful in passing from endomorphism rings to automorphism groups.

Theorem 1.4. *Let $E \to F$ be a ring isogeny. Then the induced group homomorphism $E^\times \to F^\times$ is a group isogeny. If in addition the map $E \to F$ is surjective, then so is the induced map $E^\times \to F^\times$.*

The second key result for the proof of Theorem 1.2(c) reads as follows.

Theorem 1.5. *Let B be a semisimple ring that is finitely generated as a module over its centre $\mathrm{Z}(B)$. Then $B^\times/(\mathrm{Z}(B)^\times[B^\times, B^\times])$ is an abelian group of finite exponent.*

The proof is given in Section 5 (Theorem 5.6). In fact, we prove an explicit version of Theorem 1.5. A *central simple algebra* over a field k is a ring B that is simple in the sense that it has precisely two two-sided ideals; that has centre equal to k; and that has finite dimension as a vector space over k; it is a well-known result [6, (7.22)] that, under these conditions, that dimension is a square.

Theorem 1.6. *Let k be a field, and let B be a central simple algebra over k. Let the dimension of B as a vector space over k be d^2, where d is a positive integer. Then the group $B^\times/(k^\times[B^\times, B^\times])$ is abelian of exponent dividing d.*

Our proof of Theorem 1.6 (see Theorem 5.5) makes use of Wedderburn's factorisation theorem for polynomials over division rings. Theorem 1.5 is an immediate consequence of Theorem 1.6.

In Section 6, we place ourselves in the situation of Theorem 1.2, but replacing the semisimplicity assumption on A by the condition that R be left-noetherian. We show that the category whose objects are finitely generated R-modules, and whose morphisms are certain equivalence classes of R-module correspondences, is equivalent to the category of finitely generated A-modules; Theorem 6.8 gives a precise statement. We deduce that the set of equivalence classes of commensurabilities $L \rightleftharpoons L$ forms a group that may be identified with $\mathrm{Aut}_A(Q \otimes_Z L)$ (Corollary 6.9). In Section 7 we show that $\mathrm{i} \circ \mathrm{e}$ and $\mathrm{i} \circ \mathrm{a}$ give rise to homomorphisms from this group to $\mathbb{Q}_{>0}$. If A is semisimple, then Theorem 1.5 will imply that these homomorphisms are trivial, and Theorem 1.2(c) will easily follow.

2. Isogenies and commensurabilities

This section is devoted to formal properties of isogenies and commensurabilities, and of their indices.

We begin by recalling a basic notion from category theory, for which we refer to [8, Ch. I, §11]. Let $\mathcal{C}$ be a category, and let $A \xrightarrow{f} B \xleftarrow{g} C$ be a

diagram in $\mathcal{C}$. We say that $(A \times_B C, p_0, p_1)$ is a *fibre product* of A and C over B if $A \overset{p_0}{\leftarrow} A \times_B C \overset{p_1}{\rightarrow} C$ is a diagram in $\mathcal{C}$ with the property that $fp_0 = gp_1$, and with the universal property that for any diagram $A \overset{h}{\leftarrow} D \overset{j}{\rightarrow} C$ that satisfies $fh = gj$, there exists a unique morphism $i\colon D \to A \times_B C$ such that $h = p_0 i$ and $j = p_1 i$. When a fibre product exists, it is unique up to a unique isomorphism, so in that case we may speak of *the* fibre product of A and C over B.

In the category **Grp** of groups, the fibre product of $A \overset{f}{\rightarrow} B \overset{g}{\leftarrow} C$ exists, and it is given by

$$A \times_B C = \{(a, c) \in A \times C : f(a) = g(c)\},$$

with p_0 and p_1 being the projection maps to A and C, respectively.

Throughout this section $\mathcal{C}$ will denote a category in which for every diagram $A \overset{f}{\rightarrow} B \overset{g}{\leftarrow} C$ the fibre product of A and C over B exists, equipped with a functor $\mathcal{C} \to \mathbf{Grp}$ that preserves fibre products. The main examples we have in mind are the category of groups with the identity functor, the category of rings with the functor that sends a ring to its underlying additive group, and the category of finitely generated left R-modules for a left-noetherian ring R, with the functor that sends an R-module to its underlying abelian group.

An *isogeny* in $\mathcal{C}$ is a morphism that becomes an isogeny in **Grp**. A *commensurability* in $\mathcal{C}$ is a correspondence in $\mathcal{C}$ that becomes a commensurability in **Grp**. We will often think of an isogeny $f\colon A \to B$ as a special case of a commensurability, which we will denote by c_f, namely $c_f = (A, \mathrm{id}, f)\colon A \rightleftharpoons B$.

The *index* $\mathrm{i}(f)$ of an isogeny f in $\mathcal{C}$ is defined to be the index of the image of f in **Grp**, and the index of a commensurability is defined analogously, as in the introduction.

For each of the results 2.1 – 2.6 below, it will be clear that it holds for $\mathcal{C}$ if it holds for **Grp**. We will therefore tacitly assume that $\mathcal{C} = \mathbf{Grp}$ in the proofs of those results.

Proposition 2.1. *Let A, B, C be objects in $\mathcal{C}$ and let h be the composition of two morphisms $A \overset{f}{\rightarrow} B \overset{g}{\rightarrow} C$. If two of f, g, h are isogenies, then so is the third. Moreover, we then have* $\mathrm{i}(h) = \mathrm{i}(f)\mathrm{i}(g)$.

Proof. We have an exact sequence of pointed sets

$$1 \to \ker f \to \ker h \to \ker g \to B/fA \to C/hA \to C/gB \to 1,$$

in which each map has the property that all its non-empty fibres have equal cardinality. Hence, any term that sits between two finite sets in the above sequence is itself finite. The first assertion of the proposition easily follows. Moreover, if all terms in the sequence are finite, then the alternating product of their cardinalities is 1, which proves the second assertion. □

Definition 2.2. Let $c = (X, f, g)\colon A \rightleftharpoons B$ and $d = (Y, h, j)\colon B \rightleftharpoons C$ be correspondences in $\mathcal{C}$. We define the *composition* of c with d by

$$d \circ c = (X \times_B Y, f \circ p_0, j \circ p_1)\colon A \rightleftharpoons C,$$

where p_0, p_1 are the canonical morphisms from $X \times_B Y$ to X, respectively Y.

Remark 2.3. It follows from the universal property of fibre products, and a routine diagram chase, that composition of correspondences is associative up to canonical isomorphism.

Proposition 2.4. *Let $X \xrightarrow{g} B \xleftarrow{h} Y$ be morphisms in $\mathcal{C}$, and suppose that h is an isogeny. Let $(W = X \times_B Y, p_0, p_1)$ be the fibre product of X and Y over B. Then:*

(a) *the morphism p_0 is an isogeny;*
(b) *if the image of g in* **Grp** *has finite kernel, then so does the image of p_1;*
(c) *if g is an isogeny, then so is p_1.*

Proof. We first prove part (a). Assume that h is an isogeny. We have

$$\ker p_0 = \{(1, y) \in X \times Y : h(y) = g(1) = 1\} \cong \ker h,$$

which is finite by assumption. Further, the kernel of $g\colon X \to B/hY$ is equal to $p_0 W$, so $(X : p_0 W) \leq (B : hY)$, which is also finite. So p_0 is an isogeny.

Similarly, $\ker p_1 \cong \ker g$, which proves part (b). Finally, part (c) is symmetric in X and Y, and so follows from part (a). □

Definition 2.5. A *skew correspondence* is a correspondence $c = (X, f, g)$ in which f is an isogeny.

Proposition 2.6. *If $c\colon A \rightleftharpoons B$ and $d\colon B \rightleftharpoons C$ are skew correspondences, respectively commensurabilities, then $d \circ c\colon A \rightleftharpoons C$ is a skew correspondence, respectively a commensurability. Moreover, if c and d are commensurabilities, then we have*

$$\mathrm{i}(d \circ c) = \mathrm{i}(d)\mathrm{i}(c).$$

Proof. The first two assertions follow immediately from Propositions 2.4 and 2.1. The third one follows from Proposition 2.1 and a routine diagram chase, which we leave to the reader. □

Definition 2.7. Let $c = (X, f, g)\colon A \rightleftharpoons B$ and $d = (Y, h, j)\colon A \rightleftharpoons B$ be two correspondences. We say that c and d are *equivalent* if there exists a commensurability $(W, p, q)\colon X \rightleftharpoons Y$ such that $fp = hq$ and $gp = jq$. We will call such a commensurability an *equivalence* between c and d.

Proposition 2.8. *Being equivalent in the sense of Definition 2.7 is an equivalence relation.*

Proof. The relation is clearly symmetric. Reflexivity is also clear, since an equivalence between (X, f, g) and itself is given by $(X, \mathrm{id}, \mathrm{id})\colon X \rightleftharpoons X$. Transitivity follows from Proposition 2.6. □

Definition 2.9. The *inverse* of a correspondence $c = (X, f, g)\colon A \rightleftharpoons B$ is defined to be $c^{-1} = (X, g, f)\colon B \rightleftharpoons A$.

Lemma 2.10. *Let c, $c'\colon A \rightleftharpoons B$ and $d\colon B \rightleftharpoons C$ be correspondences. Then:*

(a) *the correspondence $(d \circ c)^{-1}\colon C \rightleftharpoons A$ is equivalent to the composition $c^{-1} \circ d^{-1}$;*

(b) *if c is equivalent to c', then $c^{-1}\colon B \rightleftharpoons A$ is equivalent to $(c')^{-1}$.*

Proof. The proof is easy, and is left to the reader. □

Proposition 2.11. *Let $c, c'\colon A \rightleftharpoons B$ and $d, d'\colon B \rightleftharpoons C$ be correspondences. Suppose that c is equivalent to c', and d is equivalent to d'. Then $d \circ c$ is equivalent to $d' \circ c'$.*

Proof. Let $c = (X, f, g)$, $d = (Y, h, j)$.

First, we prove the proposition in the special case that $d' = d$, and $c' = (W, fp, gp)$, where $p\colon W \to X$ is an isogeny. Let $(X \times_B Y, p_0, p_1)$ be the fibre product of the diagram $X \xrightarrow{g} B \xleftarrow{h} Y$, and let $(W \times_B Y, p'_0, p'_1)$ be the fibre product of the diagram $W \xrightarrow{gp} B \xleftarrow{h} Y$. Thus $d \circ c = (X \times_B Y, fp_0, jp_1)$, and $d \circ c' = (W \times_B Y, fpp'_0, jp'_1)\colon A \rightleftharpoons C$. Since $gpp'_0 = hp'_1$, the universal property of fibre products guarantees the existence of a unique map $i\colon W \times_B Y \to X \times_B Y$ with the property that $pp'_0 = p_0 i$ and $p'_1 = p_1 i$:



Moreover, it is easy to see that $(W \times_B Y, p'_0, i)$ is the fibre product of the diagram $W \xrightarrow{p} X \xleftarrow{p_0} X \times_B Y$. It follows from Proposition 2.4 that i is an isogeny, which proves that $d \circ c$ is equivalent to $d \circ c'$.

Now, we prove the proposition in the special case that $d = d'$, and c' is arbitrary. Write $\sim$ for the equivalence relation between correspondences. Let $c' = (X', f', g')$, and let $(W, p, q)\colon X \rightleftharpoons X'$ be an equivalence between c and c'. Since p is an isogeny, we have $c \sim (W, fp, gp) = (W, f'q, g'q)$, and since q is an isogeny, we have $(W, f'q, g'q) \sim c'$. We deduce from the special case of the proposition that we just proved that $d \circ c \sim d \circ (W, fp, gp) \sim d \circ c'$.

Now, we prove the general case. By Lemma 2.10 and by the special case we just proved, we have $(d \circ c)^{-1} \sim c^{-1} \circ d^{-1} \sim c^{-1} \circ (d')^{-1} \sim (d' \circ c)^{-1}$. It therefore follows from Lemma 2.10(b), that $d \circ c \sim d' \circ c$. By the special case of the proposition that we have already proven, we also have $d' \circ c \sim d' \circ c'$, and the proposition follows. □

Proposition 2.12. *If c and d are two equivalent commensurabilities, then* $\mathrm{i}(c) = \mathrm{i}(d)$.

Proof. This is an immediate consequence of Proposition 2.1. □

The term "inverse" is justified by the following result.

Proposition 2.13. *Given a commensurability $c = (X, f, g)\colon A \rightleftharpoons B$, the composition $c^{-1} \circ c\colon A \rightleftharpoons A$ is equivalent to the commensurability* $(A, \mathrm{id}, \mathrm{id})$,

and the composition $c \circ c^{-1} \colon B \rightleftharpoons B$ *is equivalent to the commensurability* $(B, \mathrm{id}, \mathrm{id})$.

Proof. First, we prove the assertion on $c^{-1} \circ c$. By definition, $c^{-1} \circ c = (X \times_B X, fp_0, fp_1) \colon A \rightleftharpoons A$, where $(X \times_B X, p_0, p_1)$ is the fibre product of the diagram $X \xrightarrow{g} B \xleftarrow{g} X$. By the universal property of the fibre product, we have a unique map $i \colon X \to X \times_B X$ with the property that $p_0 i = p_1 i = \mathrm{id} \colon X \to X$.



Since g is an isogeny, it follows from Proposition 2.4 that p_0 is an isogeny. By Proposition 2.1, the morphism i is also an isogeny, so $(X, i, f) \colon X \times_B X \rightleftharpoons A$ defines an equivalence between $c^{-1} \circ c$ and $(A, \mathrm{id}, \mathrm{id})$.

The claim for $c \circ c^{-1}$ follows by applying the above argument to c^{-1} in place of c. □

Definition 2.14. We define $\mathcal{C}_{\mathrm{skew}}$ to be the category with the same objects as in $\mathcal{C}$, and where, for objects A, B, the morphisms from A to B are the equivalence classes of skew correspondences $A \rightleftharpoons B$. We also define $\mathcal{C}_{\mathrm{com}}$ to be the category with the same objects, and where the morphisms from A to B are the equivalence classes of commensurabilities $A \rightleftharpoons B$. It follows from Remark 2.3 and Propositions 2.8 and 2.11, that these are indeed categories, i.e. that composition of morphisms is well-defined and associative.

Proposition 2.13 implies that $\mathcal{C}_{\mathrm{com}}$ is a (generally large) groupoid, i.e. every morphism in $\mathcal{C}_{\mathrm{com}}$ is an isomorphism. In fact, we have the following sharper result.

Proposition 2.15. *The category* $\mathcal{C}_{\mathrm{com}}$ *is the maximal subgroupoid of* $\mathcal{C}_{\mathrm{skew}}$.

Proof. Let $c = (X, f, g) \colon A \rightleftharpoons B$ be a skew correspondence, and let $d = (Y, h, j) \colon B \rightleftharpoons A$ be a two-sided inverse in $\mathcal{C}_{\mathrm{skew}}$. So $d \circ c$ is equivalent to the commensurability $(A, \mathrm{id}, \mathrm{id}) \colon A \rightleftharpoons A$, and $c \circ d$ is equivalent to $(B, \mathrm{id}, \mathrm{id}) \colon B \rightleftharpoons B$, and in particular both compositions are commensurabilities. We wish to prove that g is then necessarily an isogeny, and for this it is enough to assume that $\mathcal{C} = \mathbf{Grp}$.

Let $(Y \times_A X, p_0, p_1)$ be the fibre product of the diagram $Y \xrightarrow{j} A \xleftarrow{f} X$, and let $(X \times_B Y, p_0', p_1')$ be the fibre product of the diagram $X \xrightarrow{g} B \xleftarrow{h} Y$. Since $c \circ d$ is a commensurability, the morphism gp_1 is an isogeny, so $(B : gX)$ is finite. Also, since $d \circ c$ is a commensurability, the morphism jp_1' is an isogeny, so $\ker p_1'$ is finite. But $\ker p_1' = \{(x, 1) \in X \times Y : g(x) = h(1) = 1\} \cong \ker g$. So g is an isogeny, as claimed. □

3. Ring isogenies

In the present section we prove that an isogeny of rings induces an isogeny of multiplicative groups.

We begin by recalling some standard ring theoretic facts, which can be found in [7].

Definition 3.1. The *Jacobson radical* of a ring E, denoted by $\mathrm{J}(E)$, is the intersection of the maximal left ideals of R.

Proposition 3.2. *Let E be a ring, and $y \in E$. Then the following are equivalent:*

(a) $y \in \mathrm{J}(E)$;
(b) *y is contained in every maximal right ideal of J;*
(c) *y annihilates every simple left E-module;*
(d) *y annihilates every simple right E-module;*
(e) $1 - xyz \in E^\times$ *for all x, $z \in E$.*

Proof. See [7, §4]. □

Lemma 3.3. *Let I be a two-sided ideal of E with $I \subset \mathrm{J}(E)$. Then the map $E^\times \to (E/I)^\times$ is surjective. Moreover, $u \in E$ is a unit if and only if $u + I$ is a unit in E/I.*

Proof. Let $u + I$ be a unit in E/I, and let $v + I$ be its inverse. Then we have uv, $vu \in 1 + I \subset 1 + \mathrm{J}(E) \subset E^\times$, so u has both a right inverse, namely $v(uv)^{-1}$, and a left inverse, namely $(vu)^{-1}v$. It follows that u is a unit in E. □

A ring is called *simple* if it has exactly two two-sided ideals. A ring E is called *semisimple* if all short exact sequences of left E-modules split.

Any semisimple ring is a finite direct product of simple rings. If E is a semisimple ring, then the opposite ring E^{opp} is also semisimple. A left-artinian ring is semisimple if and only if its Jacobson radical is 0. If E is an arbitrary ring, then $\mathrm{J}(E/\mathrm{J}(E)) = 0$. In particular, if E is left-artinian, then $E/\mathrm{J}(E)$ is semisimple. All these facts can be found in [7, §3 and §4].

Lemma 3.4. *Let E and F be rings, let $E \to F$ be a surjective ring homomorphism, and suppose that E is left-artinian. Then the induced group homomorphism $E^\times \to F^\times$ is surjective.*

Proof. First, suppose that E is semisimple. Then E can be written as the product of finitely many simple rings. Since the kernel of $E \to F$ is a two-sided ideal of E, it must be a subproduct, and F may then be identified with the complementary subproduct. Surjectivity of $E^\times \to F^\times$ is now obvious.

We pass to the general case. Denote by I the image of $\mathrm{J}(E)$ in F, which is a two-sided ideal of F. The map $E \to F$ induces a surjective ring homomorphism $E/\mathrm{J}(E) \to F/I$, where the ring $E/\mathrm{J}(E)$ is semisimple, so by the first part of the proof the induced map $(E/\mathrm{J}(E))^\times \to (F/I)^\times$ is surjective. By Lemma 3.3, the map $E^\times \to (E/\mathrm{J}(E))^\times$ is also surjective, so the map $E^\times \to (F/I)^\times$ induced by the composed ring homomorphism $E \to F/I$ is surjective as well. Now let $v \in F^\times$, and choose $u \in E^\times$ that maps to $v + I \in (F/I)^\times$. Then the image w of u in F satisfies $w \equiv v \bmod I$, so $w^{-1}v$

belongs to the image $1+I$ of $1+\mathrm{J}(E)$ in F. Let $x \in 1+\mathrm{J}(E)$ map to $w^{-1}v$. Then ux maps to $ww^{-1}v = v$, and we have $ux \in E^\times$ because $u \in E^\times$ and $x \in 1+\mathrm{J}(E) \subset E^\times$. This proves surjectivity of $E^\times \to F^\times$, as required. □

For a different proof of Lemma 3.4, see [9].

Lemma 3.5. *Let E be a ring, and let I, $J \subset E$ be two-sided ideals. Then the kernel of the natural ring homomorphism $E \to (E/I) \times (E/J)$ equals $I \cap J$, and its image is the fibre product $E/I \times_{E/(I+J)} E/J$.*

The proof is straightforward, and is left to the reader.

Lemma 3.6. *Let $E \to F$ be a surjective ring isogeny. Then the induced group homomorphism $E^\times \to F^\times$ is surjective.*

Proof. Let I be the kernel of the isogeny $E \to F$. Then I is finite, and we may identify F with E/I. Write $\operatorname{End} I$ for the endomorphism ring of the additive group of I. Let J and R, respectively, be the kernel and the image of the ring homomorphism $E \to \operatorname{End} I$ sending $a \in E$ to the map $x \mapsto ax$. Then R, being a subring of $\operatorname{End} I$, is a finite ring, J is a two-sided ideal of E, and we have a ring isomorphism $E/J \to R$. By Lemma 3.5, the combined map $E \to F \times R$ induces a ring isomorphism $\varphi\colon E/(I \cap J) \to F \times_{E/(I+J)} R$. Now we first prove that the map $(E/(I \cap J))^\times \to F^\times$ is surjective. Let $u \in F^\times$. Write v for the image of u in $(E/(I+J))^\times$. Since R is finite and hence left-artinian, by Lemma 3.4 we can choose $w \in R^\times$ mapping to $v \in (E/(I+J))^\times$. Then (u, w) belongs to $F^\times \times_{(E/(I+J))^\times} R^\times = (F \times_{E/(I+J)} R)^\times$, so $\varphi^{-1}(u, w)$ is a unit of $E/(I \cap J)$ that maps to $u \in F^\times$. This proves that $(E/(I \cap J))^\times \to F^\times$ is surjective. From $(I \cap J) \cdot (I \cap J) \subset JI = 0$ it follows that for each $x \in I \cap J$ the element $1+x$ has inverse $1-x$ and therefore belongs to $E^\times$; this implies $I \cap J \subset \mathrm{J}(E)$, so that by Lemma 3.3 the map $E^\times \to (E/(I \cap J))^\times$ is surjective. The composed map $E^\times \to F^\times$ is then surjective as well. □

Lemma 3.7. *Let E be a subring of a ring F such that the index $(F : E)$ of additive groups is finite. Then:*

(a) *the ring F has a two-sided ideal I with $I \subset E$ for which the ring F/I is finite;*
(b) *the index $(F^\times : E^\times)$ is finite.*

Proof. (a) Put $I = \{x \in F : FxF \subset E\}$. Then I is a two-sided ideal of F that is contained in E, and we proceed to show that I has finite index in F. Put $J = \{x \in F : Fx \subset E \text{ and } xF \subset E\}$. Then we have $I \subset J \subset E \subset F$. Denote by D the finite abelian group F/E, by $\operatorname{End} D$ its endomorphism ring, and by $(\operatorname{End} D)^{\mathrm{opp}}$ the ring opposite to $\operatorname{End} D$. Both of these rings are finite. The natural left and right E-module structures on D induce a ring homomorphism $E \to (\operatorname{End} D) \times (\operatorname{End} D)^{\mathrm{opp}}$ of which J is the kernel. It follows that J is a two-sided ideal of E of finite index in E. There is a well-defined group homomorphism

$$\begin{aligned} J &\to \operatorname{Hom}(D \otimes_{\mathbb{Z}} D, D) \\ x &\mapsto ((y+E) \otimes (z+E) \mapsto yxz + E) \end{aligned}$$

for $x \in J$, $y, z \in F$. Its kernel is I, and since $D \otimes_{\mathbb{Z}} D$ is finite, the group J/I is finite. Because each of F/E, E/J, J/I is finite, the ring F/I is finite. This proves (a).

(b) Let I be as in (a). Then F/I and $(F/I)^\times$ are finite, so the kernel K (say) of the natural group homomorphism $F^\times \to (F/I)^\times$ has finite index in $F^\times$. If $x \in K$, then $x^{-1} \in K$, so both x and x^{-1} are in $1+I$, which is contained in E. This proves $K \subset E^\times$, so $E^\times$ has finite index in $F^\times$ as well. This proves (b). □

Lemma 3.7 also appears in [10].

We can now prove Theorem 1.4 of the introduction, which reads as follows.

Theorem 3.8. *Let $E \to F$ be a ring isogeny. Then the induced group homomorphism $E^\times \to F^\times$ is a group isogeny. If in addition the map $E \to F$ is surjective, then so is the induced map $E^\times \to F^\times$.*

Proof. The last assertion is Lemma 3.6. For the first assertion, let I and D be the kernel, respectively the image of the map $E \to F$. Then the kernel of $E^\times \to D^\times$ is contained in $1+I$ and therefore finite, and by Lemma 3.6 the image is all of $D^\times$. Hence $E^\times \to D^\times$ is a group isogeny. Further, the inclusion map $D^\times \to F^\times$ is obviously injective, while by Lemma 3.7(b) the index of $D^\times$ in $F^\times$ is finite. Hence $D^\times \to F^\times$ is a group isogeny. By Proposition 2.1, the composed map $E^\times \to F^\times$ is also a group isogeny. □

4. Residually finite domains

This section is devoted to some properties of infinite domains all of whose proper quotients are finite.

Lemma 4.1. *Let Z be a domain such that for all non-zero $m \in Z$ the ring Z/mZ is finite, let Q be the field of fractions of Z, let V be a finite dimensional Q-vector space, and let L be a sub-Z-module of V. Then for all non-zero $m \in Z$ the Z-module L/mL is finite of order dividing $\#(Z/mZ)^{\dim_Q V}$, with equality if L is finitely generated and $Q{\cdot}L = V$.*

Proof. Let $m \in Z$ be non-zero. First suppose that L is finitely generated. Let $S \subset L$ be a finite subset that generates it as a Z-module, let $T \subset S$ be a maximal subset that is linearly independent over Q, and let $M \subset L$ be the Z-module generated by T. Then M is Z-free of rank $\#T$, so M/mM is finite of order $\#(Z/mZ)^{\#T} \le \#(Z/mZ)^{\dim_Q V}$, with equality if T is a Q-basis of V or, equivalently, if $Q{\cdot}L = V$. By maximality of T, we can, for each $s \in S$, choose a non-zero element $m_s \in Z$ such that $m_s s \in M$, and $m' = \prod_{s\in S} m_s$ is then a non-zero element of Z satisfying $m'L \subset M$. Because $M/m'M$ is finite, its subgroup $m'L/m'M$ is finite as well, and since the latter group is isomorphic to L/M and to mL/mM, we find that L/M and mL/mM are finite of the same order. The group L/mM is finite of order $\#(L/M){\cdot}\#(M/mM)$, so L/mL is also finite, of order

$$\frac{\#(L/M){\cdot}\#(M/mM)}{\#(mL/mM)} = \#(M/mM) = \#(Z/mZ)^{\#T} \le \#(Z/mZ)^{\dim_Q V},$$

with equality if $Q{\cdot}L = V$.

Passing to the general case, let U be the set of finitely generated sub-Z-modules L' of L, which is a directed partially ordered set by inclusion. Then L is the injective limit of all $L' \in U$, and L/mL is the injective limit of the modules L'/mL', all of which have order dividing $\#(Z/mZ)^{\dim_Q V}$. The

injective limit has then also order dividing the same number. This completes the proof of Lemma 4.1. □

We now prove Theorem 1.3.

Theorem 4.2. *Let Z be a domain such that for all non-zero $m \in Z$ the ring Z/mZ is finite, let Q be the field of fractions of Z, let A be a semisimple Q-algebra of finite vector space dimension over Q, and let $R \subset A$ be a sub-Z-algebra with $Q \cdot R = A$. Then R is left-noetherian and right-noetherian.*

Proof. Let I be a left ideal of R. Then $Q{\cdot}I$ is a left A-ideal, so by semisimplicity of A it is a direct summand of the left A-module A. Thus the endomorphism ring of the latter module contains an idempotent with image $Q{\cdot}I$. Since the endomorphisms of the left A-module A are the right multiplications by elements of A, it is equivalent to say that we can choose an idempotent $e \in A$ with $Ae = Q{\cdot}I$. We have $e \in Q{\cdot}I$, so we can choose a non-zero element $m \in Z$ with $me \in I$. Multiplying the chain of inclusions $Rme \subset I \subset R$ by e on the right, which when restricted to I is just the identity map, we obtain $Rme \subset I \subset Re$, where $Rme = mRe$ because m is central. By Lemma 4.1, the group Re/mRe is finite, so I/Rme is finite as well. Hence I is, as a left R-module, generated by me together with a finite set, and is therefore finitely generated. This proves that R is left-noetherian. Applying this result to A^{opp} and R^{opp}, we find that R is right-noetherian as well. □

Example 4.3. If we assume $Z \neq Q$, then the semisimplicity condition on A is actually necessary for the conclusion of Theorem 4.2 to be valid for all R. To see this, assume that A is not semisimple, or equivalently that $\mathrm{J}(A) \neq 0$, and choose a sub-Z-algebra $T \subset A$ that is finitely generated as a Z-module and satisfies $Q{\cdot}T = A$. Then the ring $R = T + \mathrm{J}(A)$ is not left-noetherian because $\mathrm{J}(A)$ is not finitely generated as a left R-ideal. If it were, then the non-zero Q-vector space $\mathrm{J}(A)/\mathrm{J}(A)^2$ would be finitely generated as a T-module and hence as a Z-module, which for $Z \neq Q$ is impossible.

Lemma 4.4. *Let Z be an infinite commutative ring. Suppose that there exists a faithful Z-module M with the property that for all non-zero $m \in Z$, M/mM is finite. Then Z is a domain.*

Proof. Let $a, b \in Z$ be non-zero. We have an exact sequence of Z-modules

$$M/bM \xrightarrow{a} M/abM \to M/aM \to 0.$$

The left and right terms are finite by assumption, so M/abM is finite. But since Z is infinite, and M is a faithful module, M is also infinite, and so $ab \neq 0$. □

The following result gives a description of the rings Z that occur in Theorem 1.2.

Theorem 4.5. *Let Z be an infinite commutative ring. Then the following assertions are equivalent:*

(a) *for each non-zero $m \in Z$, the ring Z/mZ is finite;*
(b) *the ring Z is a domain, and each non-zero prime ideal $\mathfrak{p}$ of Z is finitely generated as an ideal and has finite index in Z;*

(c) *either Z is a field, or it is a one-dimensional noetherian domain with the property that for every maximal ideal $\mathfrak{m}$ of Z the field $Z/\mathfrak{m}$ is finite.*

Proof. First we prove that (a) implies (b). From (a) it follows, by Lemma 4.4 applied to $M = Z$, that Z is a domain. Now let $\mathfrak{p}$ be a non-zero prime ideal, and let $m \in \mathfrak{p}$ be non-zero. Then we have $mZ \subset \mathfrak{p} \subset Z$, and since Z/mZ is finite, the index of $\mathfrak{p}$ in Z is finite and $\mathfrak{p}/mZ$ is finite. Hence $\mathfrak{p}$ is generated by m together with a finite set, and is therefore finitely generated.

Now we prove that (b) implies (c). By [5, Theorem 2], each commutative ring of which every prime ideal is finitely generated is noetherian. Hence (b) implies that Z is noetherian. If $\mathfrak{p}$ is a non-zero prime ideal, then $Z/\mathfrak{p}$ is a finite domain, and therefore a field. Hence each non-zero prime ideal is maximal, so Z has Krull dimension 0 or 1; in the former case it must be a field.

Finally, suppose that (c) holds. Then, we will deduce (a) by showing that for any non-zero ideal I of Z, the ring Z/I is finite. Suppose that there exists a non-zero ideal I in Z such that Z/I is infinite. Since Z is noetherian, we may, without loss of generality, assume that I is maximal among ideals with this property. So Z/I is infinite, but its quotient by any non-zero ideal is finite. It follows from Lemma 4.4, applied to $M = Z/I$, that Z/I is a domain, so I is a prime ideal of Z. It is also non-zero, so (c) implies that I is maximal, and therefore that Z/I is finite, which is a contradiction. □

5. On the units of semisimple rings

By a *division ring* we mean a ring D with the property $D^\times = D\backslash\{0\}$. If D is a division ring and n is a positive integer, then $\mathrm{M}(n, D)$ denotes the ring of n by n matrices over D. If G is a group, then G_{ab} denotes the maximal abelian quotient of G.

Lemma 5.1. *Let n be a positive integer, let D be a division ring, and for $x \in D^\times$ and $j \in \{1, \dots, n\}$, let $\delta_j(x) \in \mathrm{M}(n, D)$ be the diagonal matrix with jth entry equal to x and all other entries equal to 1. Then each map δ_j is a group homomorphism $D^\times \to \mathrm{M}(n, D)^\times$, they all induce the same group homomorphism $\bar{\delta}\colon D^\times_{\mathrm{ab}} \to \mathrm{M}(n, D)^\times_{\mathrm{ab}}$, and if $n \neq 2$ or $\#D \neq 2$ then $\bar{\delta}$ is surjective.*

Proof. It is clear that each δ_j is a group homomorphism, and that for each x all $\delta_j(x)$ are conjugate to each other, so that all δ_j induce the same map $D^\times_{\mathrm{ab}} \to \mathrm{M}(n, D)^\times_{\mathrm{ab}}$.

For i, $j \in \{1, \dots, n\}$, $i \neq j$, and $x \in D$, let $B_{ij}(x) \in \mathrm{M}(n, D)$ be the matrix obtained from the unit matrix by replacing the (i, j)-entry by x; then one has $B_{ij}(x) \in \mathrm{M}(n, D)^\times$. The subgroup of $\mathrm{M}(n, D)^\times$ generated by all $B_{ij}(x)$ is denoted by $\mathrm{SL}_n(D)$.

By [1, Chapter IV, Theorem 4.1], we have $\mathrm{M}(n, D)^\times = \mathrm{SL}_n(D)\cdot\delta_n(D^\times)$. Each $B_{ij}(x)$ is a transvection of the right D-vector space D^n in the sense of [1, Chapter IV, Definition 4.1], and if $n > 2$ or $\#D \neq 2$, then by [1, Chapter IV, Section 2] each transvection belongs to $[\mathrm{M}(n, D)^\times, \mathrm{M}(n, D)^\times]$, so that $\mathrm{SL}_n(D) \subset [\mathrm{M}(n, D)^\times, \mathrm{M}(n, D)^\times]$, and therefore

$$\mathrm{M}(n, D)^\times = [\mathrm{M}(n, D)^\times, \mathrm{M}(n, D)^\times]\cdot\delta_n(D^\times).$$

This implies that $\bar{\delta}$ is surjective. It is clearly also surjective if $n = 1$. □

Lemma 5.2. *Let n be a positive integer, let D be a division ring, and for each $x \in D^\times$ let $\iota(x) \in \mathrm{M}(n, D)^\times$ be x times the identity matrix. Then ι is a group homomorphism $D^\times \to \mathrm{M}(n, D)^\times$, and the group*

$$\mathrm{M}(n, D)^\times / (\iota(D^\times) \cdot [\mathrm{M}(n, D)^\times, \mathrm{M}(n, D)^\times])$$

is abelian of exponent dividing n.

Proof. It is clear that ι is a group homomorphism. If we have $n = \#D = 2$, then $\mathrm{M}(n, D)^\times$ is a non-abelian group of order 6, so that $\mathrm{M}(n, D)^\times_{\mathrm{ab}}$ has order 2 and the conclusion of the lemma is valid. Assume now that $n \neq 2$ or $\#D \neq 2$, so that the map $\bar{\delta}$ from Lemma 5.1 is surjective.

Denote by $\bar{\iota}\colon D^\times_{\mathrm{ab}} \to \mathrm{M}(n, D)^\times_{\mathrm{ab}}$ the map induced by ι. For each $x \in D^\times$ one has $\iota(x) = \prod_{j=1}^n \delta_j(x)$ and therefore $\bar{\iota}(x) = \bar{\delta}(x)^n$, so the surjectivity of $\bar{\delta}$ yields

$$\bar{\iota}(D^\times_{\mathrm{ab}}) = \bar{\delta}(D^\times_{\mathrm{ab}})^n = (\mathrm{M}(n, D)^\times_{\mathrm{ab}})^n,$$

and the lemma is proved. □

The following result is due to Wedderburn [11].

Theorem 5.3. *Let D be a division ring with centre $\mathrm{Z}(D)$, let $a \in D$, and let $f \in \mathrm{Z}(D)[X]$ be an irreducible polynomial with leading coefficient 1 such that $f(a) = 0$. Put $l = \deg f$. Then there exist $b_1, b_2, \ldots, b_l \in D^\times$ such that in $D[X]$ one has*

$$f = (X - b_1 a b_1^{-1}) \cdot \ldots \cdot (X - b_l a b_l^{-1}).$$

Proof. See [7, Theorem (16.9)]. □

Lemma 5.4. *Let D be a division ring that has finite vector space dimension m^2 over its centre $\mathrm{Z}(D)$, where m is a positive integer. Then the group $D^\times / (\mathrm{Z}(D)^\times \cdot [D^\times, D^\times])$ is abelian of exponent dividing m.*

Proof. Since $D^\times / (\mathrm{Z}(D)^\times \cdot [D^\times, D^\times])$ is a quotient of $D^\times_{\mathrm{ab}}$, it is an abelian group. Let $a \in D^\times$. It will suffice to show that the image $\bar{a}$ of a in the quotient $D^\times / (\mathrm{Z}(D)^\times \cdot [D^\times, D^\times])$ has order dividing m. The subfield $\mathrm{Z}(D)(a)$ of D is contained in a maximal subfield of D, and each maximal subfield of D is an extension field of $\mathrm{Z}(D)$ of degree m, by [6, (7.22)]. Hence we have $[\mathrm{Z}(D)(a) : \mathrm{Z}(D)] = l$ for some divisor l of m, and a is a zero of an irreducible polynomial $f \in \mathrm{Z}(D)[X]$ of degree l with leading coefficient 1. Using Theorem 5.3 we find $b_1, \ldots, b_l \in D^\times$ such that $b_1 a b_1^{-1} \cdot \ldots \cdot b_l a b_l^{-1} = (-1)^l f(0) \in \mathrm{Z}(D)^\times$. Mapping this identity to the abelian group $D^\times / (\mathrm{Z}(D)^\times \cdot [D^\times, D^\times])$ we obtain $\bar{a}^l = 1$, so $\bar{a}^m = 1$, as required. This proves Lemma 5.4. □

We can now prove Theorem 1.6 and deduce Theorem 1.5. We recall the statements.

Theorem 5.5. *Let k be a field, and let B be a central simple algebra over k. Let the dimension of B as a vector space over k be d^2, where d is a positive integer. Then the group $B^\times / (k^\times [B^\times, B^\times])$ is abelian of exponent dividing d.*

Proof. By [2, §14, Theorem 1], there are a positive integer n and a division ring D with $\mathrm{Z}(D) = k$ such that B is, as an algebra over k, isomorphic to $\mathrm{M}(n, D)$. Then D has finite degree m^2 over k, and $nm = d$. By Lemma 5.4, the cokernel of the natural group homomorphism $k^\times \to D^\times_{\mathrm{ab}}$ has exponent dividing m, and by Lemma 5.2 the cokernel of the natural group homomorphism $D^\times_{\mathrm{ab}} \to \mathrm{M}(n, D)^\times$ has exponent dividing n. It follows that the cokernel of the natural group homomorphism $k^\times \to \mathrm{M}(n, D)^\times_{\mathrm{ab}}$ has exponent dividing $nm = d$. □

Theorem 5.6. *Let B be a semisimple ring that is finitely generated as a module over its centre $\mathrm{Z}(B)$. Then $B^\times/(\mathrm{Z}(B)^\times[B^\times, B^\times])$ is an abelian group of finite exponent.*

Proof. In the case the semisimple ring B is simple, our hypothesis that it be finite over its centre implies that it is a central simple algebra over $\mathrm{Z}(B)$, and the assertion follows from Theorem 5.5. Generally, by [7, Chapter 1, Theorem (3.5)] the ring B is a product of finitely many semisimple rings that are simple, and the result follows from the case we just did. □

6. Skew correspondences as morphisms

In this section we formulate and prove the equivalence between categories of certain modules announced at the end of the introduction. We will need the notion of a skew correspondence (Definition 2.5), and the constructions of the categories $\mathcal{C}_{\mathrm{skew}}$ and $\mathcal{C}_{\mathrm{com}}$ (Definition 2.14).

Notation 6.1. The following assumptions will be in force throughout the present section: Z is an infinite commutative ring that satisfies the equivalent conditions of Theorem 4.5, with field of fractions Q; further, A is a Q-algebra of finite vector space dimension over Q, and R is a left-noetherian sub-Z-algebra of A with the property that $Q \cdot R = A$. By an R-module we shall always mean a left R-module. If L is a finitely generated R-module, let L_{tors} denote the set of all elements of L that have a non-zero annihilator in Z. Since the image of Z in R is central, L_{tors} is an sub-R-module.

Lemma 6.2. *Let L be a finitely generated R-module, and let U be a sub-R-module. Then U is finite if and only if it is contained in L_{tors}.*

Proof. First, we show that L_{tors} is finite. Since R is left-noetherian, L_{tors} is finitely generated as an R-module. So there exists a non-zero $m \in Z$ that annihilates L_{tors}, and L_{tors} is then a finitely generated module over the ring R/mR, which is finite by Lemma 4.1. This proves one implication.

For the converse, let $U \subset L$ be a finite sub-R-module. Then for each $x \in U$, the set $\{zx : z \in Z\}$ is finite, so that the annihilator of x in Z has finite index in Z; in particular it is non-zero, since Z is assumed to be infinite, so $x \in L_{\mathrm{tors}}$. □

Theorem 6.3. *Let L, M be two finitely generated R-modules. Then there exists a commensurability of R-modules $L \rightleftharpoons M$ if and only if there exists an isomorphism of A-modules $Q \otimes_Z L \cong Q \otimes_Z M$.*

Proof. First, suppose that we have an isomorphism $\phi\colon Q \otimes_Z L \to Q \otimes_Z M$ of A-modules. It follows from Lemma 6.2 that L is commensurable with

$\bar{L} = L/L_{\mathrm{tors}}$, and similarly for M, so by Proposition 2.6 we may assume without loss of generality that L and M are Z-torsion free. They then embed into $Q \otimes_Z L$, respectively $Q \otimes_Z M$, and by "clearing denominators", we can find $m_1, m_2 \in Z$ such that $m_1\phi(L)$ is contained in M, and $\phi(L)$ contains $m_2 M$. Since $L/m_1 L$ and $M/m_2 M$ are finite by Lemma 4.1, it follows that the correspondence $L \xleftarrow{\phi^{-1}} \phi(L) \cap M \hookrightarrow M$ is a commensurability.

Conversely, suppose that we have a commensurability $(X, f, g)\colon L \rightleftharpoons M$. Then the kernels and cokernels of f, g are finite R-modules, and so are Z-torsion modules by Lemma 6.2. They are therefore annihilated by the functor $Q \otimes_Z -$, so the maps $Q \otimes_Z f$ and $Q \otimes_Z g$ are isomorphisms. □

Lemma 6.4. *Let L, M be finitely generated R-modules, and let (X, f, g) and $(Y, h, j)\colon L \rightleftharpoons M$ be equivalent skew correspondences. Let $Q \otimes_Z f$ denote the map of A-modules $Q \otimes_Z L \to Q \otimes_Z M$ induced by f, and similarly for g, h, j. Then $(Q \otimes g) \circ (Q \otimes f)^{-1} = (Q \otimes j) \circ (Q \otimes h)^{-1}$.*

Proof. Let $(W, p, q)\colon X \rightleftharpoons Y$ be an equivalence between (X, f, g) and (Y, h, j). Since p and q are isogenies, Lemma 6.2 implies that $Q \otimes_Z p$ and $Q \otimes_Z q$ are both invertible. Moreover, we have

$$\begin{aligned} Q \otimes_Z f &= (Q \otimes_Z h) \circ (Q \otimes_Z q) \circ (Q \otimes_Z p)^{-1}, \\ Q \otimes_Z g &= (Q \otimes_Z j) \circ (Q \otimes_Z q) \circ (Q \otimes_Z p)^{-1}, \end{aligned}$$

so $(Q \otimes g) \circ (Q \otimes f)^{-1} = (Q \otimes j) \circ (Q \otimes h)^{-1}$. □

Let ${}_R\mathbf{Mod}$, respectively ${}_A\mathbf{Mod}$ denote the category of finitely generated R-modules, respectively finitely generated A-modules. By Lemma 6.4, we may define a functor $\mathcal{F}$ from ${}_R\mathbf{Mod}_{\mathrm{skew}}$ to ${}_A\mathbf{Mod}$ by sending an R-module L to the A-module $Q \otimes_Z L$, and an equivalence class of skew correspondences represented by $(X, f, g)\colon L \rightleftharpoons M$ to the map of A-modules $(Q \otimes g) \circ (Q \otimes f)^{-1}\colon Q \otimes_Z L \to Q \otimes_Z M$. The verification that $\mathcal{F}$ respects composition of morphisms, and thus does define a functor, is easy and is left to the reader.

The next three lemmas will show that the functor $\mathcal{F}$ has dense image, is full, and is faithful.

Lemma 6.5. *Any element of ${}_A\mathbf{Mod}$ is isomorphic to $\mathcal{F}(L)$ for some R-module L.*

Proof. Let V be an A-module with finite generating set B. Let L be the sub-R-module of V generated by B over R. Then the A-module $\mathcal{F}(L)$ is isomorphic to V. □

Lemma 6.6. *Let L, M be finitely generated R-modules, and let $\phi\colon \mathcal{F}(L) \to \mathcal{F}(M)$ be a morphism of A-modules. Then there exists a skew correspondence $c\colon L \rightleftharpoons M$ such that $\mathcal{F}(c) = \phi$.*

Proof. Let $\bar{L}$ be the image of L in $Q \otimes_Z L$, and let $\bar{M}$ be the image of M in $Q \otimes_Z M$. By Lemma 6.2, the natural map $f\colon L \to Q \otimes_Z L$ gives rise to a commensurability $c_L = (L, \mathrm{id}, f)\colon L \rightleftharpoons \bar{L}$, and similarly we have a commensurability $c_M\colon M \rightleftharpoons \bar{M}$. Since $\bar{L}$ and $\bar{M}$ are finitely generated as R-modules, and since $\bar{M}$ generates $Q \otimes_Z M$ over Q, we may choose a non-zero $m \in Z$ such that $m\phi(\bar{L})$ is contained in $\bar{M}$. Let g be the inclusion $m\phi(\bar{L}) \subseteq \bar{M}$, and define the correspondence $c_\phi = (\bar{L}, m, gm\phi)\colon \bar{L} \rightleftharpoons \bar{M}$. It

follows from Lemma 4.1 that c_ϕ is a skew correspondence. By Proposition 2.6, the composition $c = c_M^{-1} \circ c_\phi \circ c_L \colon L \rightleftharpoons M$ is also a skew correspondence, and it is easy to see that $\mathcal{F}(c) = \phi$. □

Lemma 6.7. *Let L, M be finitely generated R-modules, and let c, $d \colon L \rightleftharpoons M$ be two skew correspondences such that $\mathcal{F}(c) = \mathcal{F}(d)$. Then c and d are equivalent.*

Proof. Let $c = (X, f, g)$, and $d = (Y, h, j)$. We will show that c and d are equivalent by showing that the fibre product $(X \times_{L \oplus M} Y, p_0, p_1) \colon X \rightleftharpoons Y$ is a commensurability.

First, assume that the images of f, g, h, and j are Z-torsion free. Then f and g factor through X/X_{tors}, and similarly for h and j. By Lemma 6.2, the quotient maps $X \to X/X_{\mathrm{tors}}$ and $Y \to Y/Y_{\mathrm{tors}}$ are isogenies, so after replacing c and d by equivalent commensurabilities, we may assume that X and Y are Z-torsion free. It then follows from Lemma 6.2 that f, g, h, and j are injective. Since $\mathcal{F}(c) = \mathcal{F}(d)$, we have

$$(Q \otimes_Z g) \circ (Q \otimes_Z f)^{-1} = (Q \otimes_Z j) \circ (Q \otimes_Z h)^{-1},$$

and it follows that the canonical injection $X \times_{L \oplus M} Y \to X \times_L Y$ is an isomorphism. By Proposition 2.6, the fibre product $(X \times_L Y, p_0, p_1) \colon X \rightleftharpoons Y$ of the diagram $X \to M \leftarrow Y$ is a commensurability, which proves this special case of the lemma.

We now prove the general case. By applying Lemma 6.2 with $U = f(X)_{\mathrm{tors}}$, and similarly for g, h, and j, we may choose a non-zero $m \in Z$ such that the images of mf, mg, mh, and mj are Z-torsion free. It is easy to see that c is equivalent to (X, mf, mg), and d is equivalent to (Y, mh, mj). So the general case follows from the special case above. □

Theorem 6.8. *The functor $\mathcal{F} \colon {}_R\mathbf{Mod}_{\mathrm{skew}} \to {}_A\mathbf{Mod}$ is an equivalence of categories.*

Proof. The result follows by combining Lemmas 6.5, 6.6, and 6.7. □

If L is a finitely generated R-module, we let G_L denote the group of equivalence classes of commensurabilities $L \rightleftharpoons L$ under composition. It may be viewed as a full sub-groupoid of ${}_R\mathbf{Mod}_{\mathrm{com}}$ whose only object is L.

Corollary 6.9. *Let L be a finitely generated R-module. Then the map $G_L \to \mathrm{Aut}_A(Q \otimes_Z L)$, $(X, f, g) \mapsto (Q \otimes g) \circ (Q \otimes f)^{-1}$ is a group isomorphism.*

Proof. By Proposition 2.15, the category ${}_R\mathbf{Mod}_{\mathrm{com}}$ is the maximal sub-groupoid of ${}_R\mathbf{Mod}_{\mathrm{skew}}$. So Theorem 6.8 implies that the functor $\mathcal{F}$ induces an equivalence of categories from ${}_R\mathbf{Mod}_{\mathrm{com}}$ to the category whose objects are the finitely generated A-modules, and whose morphisms are the A-module isomorphisms. The corollary follows by restricting $\mathcal{F}$ to the full sub-groupoid G_L of ${}_R\mathbf{Mod}_{\mathrm{skew}}$. □

7. Automorphisms of commensurabilities

It is in the present section that we construct ring and group commensurabilities out of module commensurabilities. Here we retain the assumptions of Notation 6.1.

Let $c = (N, f, g)\colon L \rightleftharpoons M$ be a correspondence of R-modules. In the introduction we defined the endomorphism ring of c to be $\operatorname{End} c = \{(\lambda, \nu, \mu) \in (\operatorname{End} L) \times (\operatorname{End} N) \times (\operatorname{End} M) : \lambda f = f\nu,\ \mu g = g\nu\}$. We also recall the correspondence $\mathrm{e}(c) = (\operatorname{End} c, p_0, p_1)\colon \operatorname{End} L \rightleftharpoons \operatorname{End} M$, given by sending $(\lambda, \nu, \mu) \in \operatorname{End} c$ to λ and μ, respectively, and the induced correspondence of automorphism groups $\mathrm{a}(c)\colon \operatorname{Aut} L \rightleftharpoons \operatorname{Aut} M$. If $f\colon L \to M$ is an isogeny, we let c_f be the commensurability $(L, \mathrm{id}, f)\colon L \rightleftharpoons M$, as in Section 2.

Lemma 7.1. *Let $f\colon L \to M$ be an isogeny of finitely generated R-modules. Then the correspondence $\mathrm{e}(c_f)\colon \operatorname{End} L \rightleftharpoons \operatorname{End} M$ is a commensurability of rings.*

Proof. We first show that p_1 has finite kernel. We have

$$\ker p_1 = \{(\lambda, \lambda, 0) \in \operatorname{End} L \times \operatorname{End} L \times \operatorname{End} M : f\lambda = 0\} \cong \operatorname{Hom}(L, \ker f),$$

which is finite since L is finitely generated and $\ker f$ is finite by assumption.

Next, we show that the image of p_1 has finite additive index in $\operatorname{End} M$. The modules L_{tors} and $M/f(L)$ are finite, so by Lemma 6.2 there exist non-zero $m_1, m_2 \in Z$ such that m_1 annihilates L_{tors}, and m_2 annihilates $M/f(L)$. Thus, $f\colon m_1 L \to m_1 M$ is injective, and the image contains $m_1 m_2 M$, so f^{-1} defines a homomorphism $m_1 m_2 M \to m_1 L$. Given $\mu \in \operatorname{End} M$, we may therefore define $\lambda\colon L \to L, x \mapsto f^{-1}(m_1 m_2 \mu(f(x)))$, which has the property that $(\lambda, m_1 m_2 \mu) \in \operatorname{End} f$. So the image of p_1 contains $m_1 m_2 \operatorname{End} M$, which has finite additive index in $\operatorname{End} M$ by Lemma 4.1. This proves that p_1 is an isogeny.

We now show that the image of p_0 has finite additive index in $\operatorname{End} L$. Given any $\lambda \in \operatorname{End} L$, we may define $\mu\colon M \to M$, $y \mapsto f(\lambda(f^{-1}(m_1 m_2 y)))$, where m_1, m_2 are as before. We then have $(m_1 m_2 \lambda, \mu) \in \operatorname{End} f$. So the image of p_0 contains $m_1 m_2 \operatorname{End} L$, which has finite additive index in $\operatorname{End} L$ by Lemma 4.1.

Finally, we show that p_0 has finite kernel. We have

$$\begin{aligned} \ker p_0 &= \{(0, 0, \mu) \in \operatorname{End} L \times \operatorname{End} L \times \operatorname{End} M : \mu f = 0\} \\ &\cong \operatorname{Hom}(M/f(L), M) \cong \operatorname{Hom}(M/f(L), M_{\mathrm{tors}}), \end{aligned}$$

where the last isomorphism follows from Lemma 6.2 and the assumption that $M/f(L)$ is finite. Invoking Lemma 6.2 again, it follows that $\ker p_0$ is finite, so p_0 is an isogeny. □

Theorem 7.2. *Let L, M be finitely generated R-modules. Then for any commensurability $c = (X, f, g)\colon L \rightleftharpoons M$, the correspondence $\mathrm{e}(c)\colon \operatorname{End} L \rightleftharpoons \operatorname{End} M$ is a ring commensurability, and the induced correspondence $\mathrm{a}(c)\colon \operatorname{Aut} L \rightleftharpoons \operatorname{Aut} M$ is a group commensurability.*

Proof. The correspondence $\mathrm{e}(c)$ is canonically isomorphic to the composition of $\mathrm{e}(c_f)^{-1}\colon \operatorname{End} L \rightleftharpoons \operatorname{End} X$ with $\mathrm{e}(c_g)\colon \operatorname{End} X \rightleftharpoons \operatorname{End} M$. The correspondences $\mathrm{e}(c_f)$ and $\mathrm{e}(c_g)$ are commensurabilities by Lemma 7.1, so $\mathrm{e}(c)$ is a commensurability by Proposition 2.6. The assertion on $\mathrm{a}(c)$ follows from Theorem 3.8 by passing to the unit groups. □

Theorem 7.3. *Let $c\colon L \rightleftharpoons M$, $d\colon M \rightleftharpoons N$ be commensurabilities of R-modules. Then:*

(a) *the ring commensurability* $\mathrm{e}(d \circ c)\colon \operatorname{End} L \rightleftharpoons \operatorname{End} N$ *is equivalent (see Definition 2.7) to the composition of ring commensurabilities* $\mathrm{e}(d) \circ \mathrm{e}(c)$*, and the group commensurability* $\mathrm{a}(d \circ c)$ *is equivalent to the composition* $\mathrm{a}(d) \circ \mathrm{a}(c)$*;*

(b) *we have*

$$\begin{aligned} \mathrm{i}(\mathrm{e}(d \circ c)) &= \mathrm{i}(\mathrm{e}(d))\mathrm{i}(\mathrm{e}(c)), \\ \mathrm{i}(\mathrm{a}(d \circ c)) &= \mathrm{i}(\mathrm{a}(d))\mathrm{i}(\mathrm{a}(c)). \end{aligned}$$

Proof. (a) Write $c = (X, f, g)\colon L \rightleftharpoons M$, $d = (Y, h, j)\colon M \rightleftharpoons N$. We claim that there is an isogeny

$$i\colon \operatorname{End} c \times_{\operatorname{End} M} \operatorname{End} d \to \operatorname{End}(d \circ c)$$

that makes the following diagram of endomorphism rings commute:



where all unlabelled morphisms are the ones defined in the introduction.

An element of $\operatorname{End} c \times_{\operatorname{End} M} \operatorname{End} d$ is a pair of triples

$$\begin{aligned} &((\lambda, \xi, \mu), (\mu', \upsilon, \nu)), \\ &\lambda \in \operatorname{End} L, \xi \in \operatorname{End} X, \mu, \mu' \in \operatorname{End} M, \upsilon \in \operatorname{End} Y, \nu \in \operatorname{End} N, \end{aligned}$$

satisfying $\lambda f = f\xi$, $\mu g = g\xi$, $\mu' h = h\upsilon$, $\nu j = j\upsilon$, and the fibre product condition in fact demands that $\mu = \mu'$.

An element of $\operatorname{End}(d \circ c)$ is a triple $(\lambda', \zeta', \nu') \in \operatorname{End} L \times \operatorname{End}(X \times_M Y) \times \operatorname{End} N$ satisfying $\lambda' f p_0 = \zeta' f p_0$, $\nu' j p_1 = j p_1 \zeta'$, where p_0, p_1 are the canonical projection maps from $X \times_M Y$ to X, respectively Y. Define

$$\begin{array}{rccc} i\colon & \operatorname{End} c \times_{\operatorname{End} M} \operatorname{End} d & \to & \operatorname{End}(d \circ c) \\ & ((\lambda, \xi, \mu), (\mu, \upsilon, \nu)) & \mapsto & (\lambda, (\xi, \upsilon), \nu). \end{array}$$

A routine diagram chase, which we leave to the reader, shows that the image of i is indeed contained in $\operatorname{End}(d \circ c)$.

To see that this definition of i makes the above diagram of endomorphism rings commute is also routine, and will also be omitted. It remains to check that i is an isogeny. The correspondence $\mathrm{e}(d) \circ \mathrm{e}(c)\colon \operatorname{End} L \rightleftharpoons \operatorname{End} N$ consists of $\operatorname{End} c \times_{\operatorname{End} M} \operatorname{End} d$, together with the maps to $\operatorname{End} L$ and $\operatorname{End} N$. By Theorem 7.2, the correspondences $\mathrm{e}(c)$ and $\mathrm{e}(d)$ are commensurabilities, so by Proposition 2.6 the correspondence $\mathrm{e}(d) \circ \mathrm{e}(c)$ is a commensurability. In particular, the morphism $\operatorname{End} c \times_{\operatorname{End} M} \operatorname{End} d \to \operatorname{End} L$ is an isogeny. Also,

$\mathrm{End}(d \circ c) \to \mathrm{End}\, L$ is an isogeny by Theorem 7.2. The fact that i is an isogeny therefore follows from Proposition 2.1. This proves our claim.

The isogeny i defines an equivalence between $\mathrm{e}(d \circ c)$ and $\mathrm{e}(d) \circ \mathrm{e}(c)$. This proves part (a) for endomorphism rings. By passing to the unit groups and applying Theorem 3.8 to the isogeny i, we also obtain part (a) for automorphism groups.

Part (b) immediately follows from part (a) by Propositions 2.12 and 2.6. □

Proposition 7.4. *Let L, M be finitely generated R-modules, and let c, $d\colon L \rightleftharpoons M$ be two commensurabilities. If c is equivalent to d, then $\mathrm{e}(c)$ is equivalent to $\mathrm{e}(d)$, and $\mathrm{a}(c)$ is equivalent to $\mathrm{a}(d)$.*

Proof. Let $c = (X, f, g)$ and $d = (Y, h, j)$. First, assume that an equivalence between c and d is given by an isogeny $p\colon Y \to X$, so that we have the following commutative diagram:



As before, write $c_f = (X, \mathrm{id}, f)\colon X \rightleftharpoons L$, and define c_g, c_p similarly. Then d is canonically isomorphic to $(c_g \circ c_p) \circ (c_p^{-1} \circ c_f^{-1})$. By Theorem 7.3, the commensurability $\mathrm{e}(d)$ is equivalent to $\mathrm{e}(c_g) \circ \mathrm{e}(c_p) \circ \mathrm{e}(c_p)^{-1} \circ \mathrm{e}(c_f^{-1})$. By Proposition 2.13, the composition $\mathrm{e}(c_p) \circ \mathrm{e}(c_p)^{-1}$ is equivalent to $(\mathrm{End}\, X, \mathrm{id}, \mathrm{id})\colon \mathrm{End}\, X \rightleftharpoons \mathrm{End}\, X$. So by Proposition 2.11 the commensurability $\mathrm{e}(c_g) \circ \mathrm{e}(c_p) \circ \mathrm{e}(c_p)^{-1} \circ \mathrm{e}(c_f^{-1})$ is equivalent to $\mathrm{e}(c_g) \circ (\mathrm{End}\, X, \mathrm{id}, \mathrm{id}) \circ \mathrm{e}(c_f^{-1})$, which is canonically isomorphic to $\mathrm{e}(c_g) \circ \mathrm{e}(c_f^{-1})$. Applying Theorem 7.3 again, we find that $\mathrm{e}(c_g) \circ \mathrm{e}(c_f^{-1})$ is equivalent to $\mathrm{e}(c_g \circ c_f^{-1})$. Finally, $c_g \circ c_f^{-1}$ is canonically isomorphic to c, and the special case of the proposition follows.

Passing to the general case, let $(W, p, q)\colon X \rightleftharpoons Y$ be an equivalence between c and d. Since p is an isogeny, c is equivalent to (W, fp, gp), and since q is an isogeny, d is equivalent to $(W, hq, jq) = (W, fp, gp)$. The result therefore follows from the special case we just did. □

Let **Rng** denote the category of rings, and **Grp** the category of groups. Theorem 7.3 and Proposition 7.4 imply that there is a functor from ${}_R\mathbf{Mod}_{\mathrm{com}}$ to $\mathbf{Rng}_{\mathrm{com}}$ that takes an R-module L to the ring $\mathrm{End}\, L$, and an equivalence class of R-module commensurabilities, represented by a commensurability c, to the equivalence class of ring commensurabilities represented by $\mathrm{e}(c)$. Further, Theorem 3.8 shows that we have the functors $^+$ and $^\times$ from $\mathbf{Rng}_{\mathrm{com}}$ to $\mathbf{Grp}_{\mathrm{com}}$ which take a ring to the additive, respectively multiplicative group of the ring. Finally, Propositions 2.12 and 2.6 imply that we have the functor i from $\mathbf{Grp}_{\mathrm{com}}$ to the group $\mathbb{Q}_{>0}$, thought of as a groupoid with one object. To summarise, we have the functors of groupoids

$$(7.5) \qquad {}_R\mathbf{Mod}_{\mathrm{com}} \xrightarrow{\mathrm{End}} \mathbf{Rng}_{\mathrm{com}} \underset{\times}{\overset{+}{\rightrightarrows}} \mathbf{Grp}_{\mathrm{com}} \xrightarrow{\mathrm{i}} \mathbb{Q}_{>0}.$$

Let L be a finitely generated R-module, and let V denote the A-module $Q \otimes_Z L$. The isomorphism of Corollary 6.9 and the functors (7.5) then induce

group homomorphisms

$$\begin{array}{rccl} \operatorname{Aut}_A V \cong & G_L & \to & \mathbb{Q}_{>0}, \\ & c & \mapsto & \mathrm{i}(\mathrm{e}(c)) \quad \text{and} \\ & c & \mapsto & \mathrm{i}(\mathrm{a}(c)). \end{array} \tag{7.6}$$

Lemma 7.7. *Let L be a finitely generated R-module, write $\bar{L} = L/L_{\mathrm{tors}}$, and let $f\colon L \to \bar{L}$ denote the quotient map. Then the isomorphism in ${}_R\mathbf{Mod}_{\mathrm{com}}$ given by the commensurability $(L, \mathrm{id}, f)\colon L \rightleftharpoons \bar{L}$ induces an isomorphism $G_L \to G_{\bar{L}}$ that commutes with the maps $G_L \to \mathbb{Q}_{>0}$ and $G_{\bar{L}} \to \mathbb{Q}_{>0}$ defined in (7.6).*

Proof. Let t denote the commensurability $L \xleftarrow{\mathrm{id}} L \to \bar{L}$. Then the isomorphism $G_L \to G_{\bar{L}}$ is given by composition on the right with t and on the left with t^{-1}. It easily follows from Theorem 7.3, Proposition 2.6 and Proposition 2.12 that this isomorphism commutes with the maps (7.6). □

Proposition 7.8. *Let L be a finitely generated R-module, and denote the A-module $Q \otimes_Z L$ by V. Let α be an element of $\mathrm{Z}(\operatorname{End}_A V)^\times \subseteq \operatorname{Aut}_A V \cong G_L$. Then its image in $\mathbf{Rng}_{\mathrm{com}}$ under the first functor of (7.5) is the identity morphism on $\operatorname{End} L$.*

Proof. By Lemma 7.7, we may assume that L is Z-torsion free. Thus, L injects into $V = Q \otimes_Z L$. For any sub-R-module U of V, write $E_U = \{\phi \in \operatorname{End}_A V : \phi U \subset U\}$. Then the injection $L \hookrightarrow V$ induces a map $\operatorname{End}_R L \to \operatorname{End}_A V$, which is injective and whose image is exactly E_L.

Let $\alpha \in \operatorname{Aut}_A V$ be arbitrary. Then the isomorphism $\operatorname{Aut}_A V \cong G_L$ identifies α with the equivalence class of commensurabilities represented by $c = (L \cap \alpha^{-1}L, i, \alpha)\colon L \rightleftharpoons L$, where $i\colon L \cap \alpha^{-1}L \to L$ is the inclusion map. We have

$$\begin{aligned} \operatorname{End} c = \{&(\lambda_0, \lambda_1) \in \operatorname{End}_A V \times \operatorname{End}_A V : \\ &\lambda_0 \in E_L \cap E_{\alpha^{-1}L}, \lambda_1 \in E_{\alpha L} \cap E_L, \lambda_0 = \alpha^{-1}\lambda_1\alpha\}. \end{aligned}$$

The commensurability $\mathrm{e}(c)$ is then of the form $(\operatorname{End} c, p_0, p_1)\colon \operatorname{End} L \rightleftharpoons \operatorname{End} L$, where $p_0\colon (\lambda_0, \lambda_1) \mapsto \lambda_0$, and $p_1\colon (\lambda_0, \lambda_1) \mapsto \lambda_1 = \alpha\lambda_0\alpha^{-1}$.

It follows that if α is an element of $\mathrm{Z}(\operatorname{End}_A V)^\times$, then p_0 and p_1 are equal. In this case, the commensurability $(\operatorname{End} c, \mathrm{id}, p_0)\colon \operatorname{End} c \rightleftharpoons \operatorname{End} L$ defines an equivalence between $\mathrm{e}(c)$ and $(\operatorname{End} L, \mathrm{id}, \mathrm{id})\colon \operatorname{End} L \rightleftharpoons \operatorname{End} L$, the identity morphism on $\operatorname{End} L$ in $\mathbf{Rng}_{\mathrm{com}}$. □

The following result is an immediate consequence of Proposition 7.8.

Corollary 7.9. *The two group homomorphisms $\operatorname{Aut}_A V \to \mathbb{Q}_{>0}$ of (7.6) factor through $\operatorname{Aut}_A V/\mathrm{Z}(\operatorname{End}_A V)^\times$.*

Remark 7.10. The computation in the proof of Proposition 7.8 shows that the group homomorphism $\mathrm{i} \circ \mathrm{e}\colon \operatorname{Aut}_A V \to \mathbb{Q}_{>0}$ is given by

$$\alpha \mapsto \frac{(E_L : E_{\alpha L} \cap E_L)}{(E_L : E_L \cap E_{\alpha^{-1}L})},$$

and analogously for $\mathrm{i} \circ \mathrm{a}$.

8. The case of semisimple algebras

In this section, we will prove our main results. We begin with Theorem 1.2. We recall the statement.

Theorem 8.1. *Let Z be an infinite domain such that for all non-zero $m \in Z$ the ring Z/mZ is finite, let Q be the field of fractions of Z, let A be a semisimple Q-algebra of finite vector space dimension over Q, let $R \subset A$ be a sub-Z-algebra with $Q \cdot R = A$, and let L, M be finitely generated R-modules. Then:*

(a) *there is an R-module commensurability $L \rightleftharpoons M$ if and only if the A-modules $Q \otimes_Z L$ and $Q \otimes_Z M$ are isomorphic;*
(b) *if $c\colon L \rightleftharpoons M$ is an R-module commensurability, then $\mathrm{e}(c)\colon \operatorname{End} L \rightleftharpoons \operatorname{End} M$ is a ring commensurability, and $\mathrm{a}(c)\colon \operatorname{Aut} L \rightleftharpoons \operatorname{Aut} M$ is a group commensurability;*
(c) *if c, $c'\colon L \rightleftharpoons M$ are R-module commensurabilities, then one has*

$$\mathrm{i}(\mathrm{e}(c)) = \mathrm{i}(\mathrm{e}(c')), \qquad \mathrm{i}(\mathrm{a}(c)) = \mathrm{i}(\mathrm{a}(c')).$$

Proof. By Theorem 4.2, the ring R is left-noetherian, so the assumptions of Notation 6.1 are satisfied. Parts (a) and (b) of the theorem therefore follow from Theorems 6.3 and 7.2, respectively.

We now prove part (c). Let $c, c'\colon L \rightleftharpoons M$ be R-module commensurabilities. By Theorem 7.3, the assertion of part (c) is equivalent to the statement that

$$\mathrm{i}(\mathrm{e}(c^{-1} \circ c')) = \mathrm{i}(\mathrm{a}(c^{-1} \circ c')) = 1.$$

So we may, without loss of generality, assume that $L = M$, and it will suffice to show that the homomorphisms

$$\mathrm{i} \circ \mathrm{e}, \mathrm{i} \circ \mathrm{a}\colon \operatorname{Aut}_A V \cong G_L \to \mathbb{Q}_{>0}$$

defined in (7.6) are trivial. Here V denotes the A-module $Q \otimes_Z L$.

Let B denote the Q-algebra $\operatorname{End}_A V$, so that $G_L = B^\times$. Since $\mathbb{Q}_{>0}$ is abelian, both homomorphisms $\mathrm{i} \circ \mathrm{e}$ and $\mathrm{i} \circ \mathrm{a}$ factor through $B^\times/[B^\times, B^\times]$. By Corollary 7.9, they also factor through $B^\times/\mathrm{Z}(B)^\times$. Since A is a semisimple ring, and since V is a finitely generated A-module, it follows that V is a finite direct sum of simple modules, so by Schur's lemma B is a direct product of matrix rings over division rings, and in particular a semisimple ring. By Theorem 5.6, the quotient $B^\times/(\mathrm{Z}(B)^\times[B^\times, B^\times])$ is an abelian group of finite exponent. Since $\mathbb{Q}_{>0}$ is torsion-free, any homomorphism $B^\times/(\mathrm{Z}(B)^\times[B^\times, B^\times]) \to \mathbb{Q}_{>0}$ must be trivial. □

Example 8.2. The following example demonstrates that if we replace the semisimplicity assumption on A by the condition that R be left-noetherian, then the conclusion of Theorem 1.2(c) need no longer hold.

Let $R = \left(\begin{smallmatrix} \mathbb{Z} & \mathbb{Z} \\ 0 & \mathbb{Z} \end{smallmatrix}\right)$, and $A = \mathbb{Q} \otimes_{\mathbb{Z}} R$. Let L be a free R-module of rank 1, set $V = \mathbb{Q} \otimes_{\mathbb{Z}} L$, and $B = \operatorname{End}_A V$. We have $\operatorname{End} L \cong R^{\mathrm{opp}}$, and similarly

$$B^\times \cong (A^{\mathrm{opp}})^\times \cong \left(\begin{smallmatrix} \mathbb{Q}^\times & 0 \\ \mathbb{Q} & \mathbb{Q}^\times \end{smallmatrix}\right). \tag{8.3}$$

Recall from equation (7.6), that $\mathrm{i} \circ \mathrm{e}$ defines a group homomorphism from $B^\times$ to $\mathbb{Q}_{>0}$, which factors through $B^\times/(\mathrm{Z}(B)^\times \cdot [B^\times, B^\times])$. The map $\left(\begin{smallmatrix} a & 0 \\ b & c \end{smallmatrix}\right) \mapsto c/a$ defines an isomorphism of this quotient with $\mathbb{Q}^\times$. For $\alpha = \left(\begin{smallmatrix} 1 & 0 \\ 0 & c \end{smallmatrix}\right)$, one

easily computes, using Remark 7.10, that $\mathrm{i} \circ \mathrm{e}(\alpha) = \mathrm{i} \circ \mathrm{a}(\alpha) = |c|$. It follows that both $\mathrm{i} \circ \mathrm{e}$ and $\mathrm{i} \circ \mathrm{a}$ map $\left(\begin{smallmatrix} a & 0 \\ b & c \end{smallmatrix}\right)$ to $|c/a|$, and are therefore far from trivial.

We now deduce Theorem 1.1.

Theorem 8.4. *Let G be a finite group, let V be a finitely generated $\mathbb{Q}[G]$-module, and put $\mathcal{S} = \{L : L$ is a finitely generated $\mathbb{Z}[G]$-module with $\mathbb{Q} \otimes_{\mathbb{Z}} L \cong V$ as $\mathbb{Q}[G]$-modules$\}$. Then there exists a unique function $\mathrm{ia} \colon \mathcal{S} \times \mathcal{S} \to \mathbb{Q}_{>0}$ such that*

(a) *if L, L', M, $M' \in \mathcal{S}$ and $L \cong L'$, $M \cong M'$, then $\mathrm{ia}(L, M) = \mathrm{ia}(L', M')$;*
(b) *if L, M, $N \in \mathcal{S}$, then $\mathrm{ia}(L, M) \cdot \mathrm{ia}(M, N) = \mathrm{ia}(L, N)$;*
(c) *if $M \in \mathcal{S}$, and $L \subset M$ is a submodule of finite index, then with $H = \{\sigma \in \operatorname{Aut} M : \sigma L = L\}$ and $\rho \colon H \to \operatorname{Aut} L$ mapping $\sigma \in H$ to $\sigma|L$, one has*

$$\mathrm{ia}(L, M) = \frac{(\operatorname{Aut} M : H) \cdot \# \ker \rho}{(\operatorname{Aut} L : \rho H)}.$$

Proof. Existence immediately follows from Theorem 8.1. In particular, property (c) follows by taking the commensurability $c = (L, \mathrm{id}, i) \colon L \rightleftharpoons M$, where $i \colon L \to M$ is the inclusion map, and noticing that H is precisely $\operatorname{Aut} c$.

To show uniqueness, note that the conditions of the theorem imply that the function ia, if it exists, is uniquely determined by its values on $\mathbb{Z}$-free modules. Indeed, if m_1 and m_2 are the exponents of the $\mathbb{Z}$-torsion submodule of L, respectively of M, then condition (b) requires that

$$\mathrm{ia}(m_1 L, L)\, \mathrm{ia}(L, M) = \mathrm{ia}(m_1 L, m_2 M)\, \mathrm{ia}(m_2 M, M).$$

Condition (c) determines the values of $\mathrm{ia}(m_1 L, L)$ and $\mathrm{ia}(m_2 M, M)$, so $\mathrm{ia}(L, M)$ is determined by $\mathrm{ia}(m_1 L, m_2 M)$. Clearly, the modules $m_1 L$ and $m_2 M$ are both $\mathbb{Z}$-free.

But if L, M are $\mathbb{Z}$-free, and $\mathbb{Q} \otimes_{\mathbb{Z}} L \cong \mathbb{Q} \otimes_{\mathbb{Z}} M$, then there exists an embedding $L \hookrightarrow M$ with finite index, so that $\mathrm{ia}(L, M)$ is determined by conditions (a) and (c). □

The first interesting case of Theorem 1.1 is already when G is the trivial group, so that finitely generated $\mathbb{Z}[G]$-modules are just finitely generated abelian groups.

Proposition 8.5. *Let L, M be finitely generated abelian groups. Then:*

(a) *there exists a commensurability $L \rightleftharpoons M$ if and only if L and M have the same rank;*
(b) *if $L \cong \mathbb{Z}^n \oplus L_0$ and $M \cong \mathbb{Z}^n \oplus M_0$, where L_0 and M_0 are finite abelian groups, then*

$$\mathrm{ia}(L, M) = \frac{(\# M_0)^n \cdot \# \operatorname{Aut} M_0}{(\# L_0)^n \cdot \# \operatorname{Aut} L_0}.$$

Proof. Part (a) immediately follows from Theorem 1.2(a).

We now prove part (b). First we compute $\mathrm{ia}(\mathbb{Z}^n, L)$. The split exact sequence

$$0 \to L_0 \to L \xrightarrow{f} \mathbb{Z}^n \to 0$$

induces a surjective map

$$\operatorname{Aut} L \to \operatorname{Aut} L_0 \oplus \operatorname{Aut} \mathbb{Z}^n,$$

whose kernel is easily seen to be canonically isomorphic to $\mathrm{Hom}(\mathbb{Z}^n, L_0)$. It follows that if c is the commensurability $(L, f, \mathrm{id}) : \mathbb{Z}^n \rightleftharpoons L$, then the map $\mathrm{Aut}\, c \to \mathrm{Aut}\, L$ is an isomorphism, while the map $\mathrm{Aut}\, c \to \mathrm{Aut}\, \mathbb{Z}^n$ is onto, with kernel of cardinality $\#\mathrm{Hom}(\mathbb{Z}^n, L_0) \cdot \#\mathrm{Aut}\, L_0 = (\#L_0)^n \cdot \#\mathrm{Aut}\, L_0$. Hence $\mathrm{ia}(\mathbb{Z}^n, L) = \mathrm{i}(\mathrm{a}(c)) = (\#L_0)^n \cdot \#\mathrm{Aut}\, L_0$.

It follows from the above computation that

$$\begin{aligned} \mathrm{ia}(L, M) &= \frac{\mathrm{ia}(\mathbb{Z}^n, M)}{\mathrm{ia}(\mathbb{Z}^n, L)} \\ &= \frac{(\#M_0)^n \cdot \#\mathrm{Aut}\, M_0}{(\#L_0)^n \cdot \#\mathrm{Aut}\, L_0}, \end{aligned}$$

as claimed. □

References

[1] E. Artin, Geometric Algebra, Interscience Publishers Inc., New York (1957).

[2] N. Bourbaki, Éléments de mathématique, Algèbre, Chapitre 8, Modules et anneaux semi-simples, Appendice 2.

[3] H. Cohen and H. W. Lenstra, Jr., Heuristics on class groups of number fields, Number theory, Noordwijkerhout 1983, Lecture Notes in Math. 1068, Springer, Berlin (1984), 33–62.

[4] H. Cohen and J. Martinet, Étude heuristique des groupes de classes des corps de nombres, J. Reine Angew. Math. 404 (1990), 39–76.

[5] I. S. Cohen, Commutative rings with restricted minimum condition, Duke Math. J. 17, (1950), 27–42.

[6] C. W. Curtis and I. Reiner, Methods of Representation Theory with Applications to Finite Groups and Orders, Vol. 1, John Wiley & Sons, New York (1981).

[7] T. Y. Lam, A First Course in Noncommutative Rings, GTM 131, Springer, New York (2001).

[8] S. Lang, Algebra, GTM 211, Springer, New York (2002).

[9] H. W. Lenstra, A normal basis theorem for infinite Galois extensions, Indag. Math. 88 (1985), 221–228.

[10] J. Lewin, Subrings of finite index in finitely generated rings, J. Algebra 5 (1967), 84–88.

[11] J. H. M. Wedderburn, On division algebras, Trans. Amer. Math. Soc. 22 (1921), 129–135.